Allergy Action Plan Student's Name: D.O.B:

1 1211111111111111111111111111111111111	D.O.B:Teacher:	1 .
ALLE	Child's Picture	
	RGY TO: atic Yes* No *Fligher risk for severe reaction	Here
	◆ STEP 1: TREATMENT ◆	
Symp	Give Checked Medicar (To be determined by physician aut	hon**:
" If a	food allergen has been ingested, but no symptoms:	l Antihistamine
* Mo	uth Itching tingling or swelling of line toward	Antihistamine
- Skir	Hives, itchy rash, swelling of the face or extraorities	Antihistamine
* Gut	Nausca, abdominal crames vomiting disables	Antihistamine
* Thro	Out! Tightening of throat boarseness bashing and	Antihistamine
• Lun	Shortness of breath repetitive conching whencing	Antihistamine
- Hear	Thready pulse, low blood pressure fainting note him	Antihistamine
* Othe	The Designation of the Prince	Antihistamine
 If rea 	iction is progressing (several of the above areas affected), give	Antihistamine
The severi	ry of symptoms can quickly change. †Potentially life-threatening.	- Middingtalining
	rine: inject intramuscularly (circle one) EpiPen® EpiPen® Jr. Twinject™ 0.3 mg Twin rse side for instructions) unine: give	
	medication/dose/route	1944
Other: gi	Ve	
	medication/dose/route	
	♦ <u>STEP 2: EMERGENCY CALLS</u> ♦	
1. Call 91 may be	I (or Rescue Squad:). State that an allergic reaction has been treated needed.	, and additional epinephrine
2. Dr		
3. Emerger Namc/Relat	acy contacts;	
a	• • • • • • • • • • • • • • • • • • • •	
		17/00/04
	1.)	NOTE 11 and 12 a
	1.)	
EVEN IF PAF	CENTAGUARDIAN CANNOT BE REACHED, DO NOT HESITATE TO MEDICATE OR TAKE CHIED TO M	THE TAIL TO STATE STATE OF THE
Parent/Guard	ray pighatha	
	Date	

 $\mathbf{p}_{\mathbf{a}}$ Doctor's Signature___ (Required) Date__